## Missouri School Immunization Requirements 2009-2010

- All students must present documentation of month, day, and year of each immunization before they attend school.
- All immunizations must be up-to-date before students are permitted to attend classes.
- The Advisory Committee on Immunization Practices (ACIP) allows a 4-day grace period, so students in all grade levels may receive immunizations up to 4 days before they are due.
- For children beginning kindergarten during or after the 2003-04 school year, required immunizations should be administered according to the current ACIP Schedule (http://www.cdc.gov/vaccines/recs/schedules/child-schedule.htm).
- To remain in school, students "in progress" must have an Imm.P.14 form on file and must receive immunizations as soon as they become due.
- Religious (Imm.P.11A) and medical (Imm.P.12) exemptions are allowed. The appropriate exemption card must be on file.

Grades	Immunizations	
K-4	4+ DTaP/DTP/DT/Td <sup>1</sup> 3+ polio <sup>2</sup> 2 MMR (measles, mumps, rubella)	3+ hepatitis B  1 varicella (chickenpox) or proof of disease <sup>3</sup>
5-6	4+ DTaP/DTP/DT/Td <sup>1</sup> 3+ polio <sup>2</sup>	2 MMR (measles, mumps, rubella) 3+ hepatitis B
7-10	4 DTaP/DTP/DT/Td <sup>1, 4</sup> 3+ polio <sup>2</sup>	2 measles, 1 mumps, 1 rubella, 3+ hepatitis B <sup>5</sup>
11-12	3+ DTaP/DTP/DT/Td <sup>4</sup> 3+ polio <sup>2</sup>	2 measles, 1 mumps, 1 rubella, 3+ hepatitis B <sup>5</sup>

- 1. Last dose on or after fourth (4th) birthday and last dose of pediatric pertussis before seventh (7th) birthday. **Maximum needed**: six (6) doses.
- 2. Last dose on or after fourth (4th) birthday. If a combination of IPV/OPV is received, four (4) doses are required. **Maximum needed**: four (4) doses.
- 3. A statement signed by parent/guardian or physician indicating the month and year the child had chickenpox disease.
- 4. The ACIP and the Missouri Department of Health and Senior Services recommend Tdap, which contains pertussis vaccine, at age 11-12 years for those who have completed the recommended childhood DTP/DTaP vaccination series and have not received a Td booster dose. However, pertussis vaccine is not required for adolescents for school attendance at this time. A Td booster is required ten (10) years after the last dose of DTaP, DTP, DT, or Td.
- 5. Students who are 11-15 years of age may receive two doses adult formulation Merck Recombivax Hepatitis B vaccine on the following schedule:

Dose 1, initial visit

Dose 2, 4-6 months after Dose 1

